

Resource Summary Report

Generated by [ASWG](#) on Aug 10, 2026

Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, DyLight™ 680

RRID:AB_2556622

Type: Antibody

Proper Citation

Thermo Fisher Scientific Cat# SA5-10042, RRID:AB_2556622

Antibody Information

URL: http://antibodyregistry.org/AB_2556622

Proper Citation: Thermo Fisher Scientific Cat# SA5-10042, RRID:AB_2556622

Target Antigen: Rabbit IgG (H+L)

Host Organism: donkey

Clonality: polyclonal secondary

Comments: Applications: Flow (1:50 - 1:200), ICC/IF (1:50-1:500), IHC (1:50-1:500), IP (Assay-dependent), WB (1:1,000-1:20,000)

Antibody Name: Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, DyLight™ 680

Description: This polyclonal secondary targets Rabbit IgG (H+L)

Target Organism: rabbit

Antibody ID: AB_2556622

Vendor: Thermo Fisher Scientific

Catalog Number: SA5-10042

Record Creation Time: 20250819T223715+0000

Record Last Update: 20250820T024623+0000

Ratings and Alerts

No rating or validation information has been found for Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, DyLight™ 680.

No alerts have been found for Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, DyLight™ 680.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 7 mentions in open access literature.

Listed below are recent publications. The full list is available at [ASWG](#) .

Lesire L, et al. (2026) Enhancing antitumour response to proteasome inhibitors with inhibitors of insulin-degrading enzyme, a new molecular vulnerability in multiple myeloma. British journal of pharmacology, 183(12), 3355.

Mikolajewicz N, et al. (2026) Reactive oligodendrocytes promote glioblastoma progression through CCL5/CCR5-mediated glioma stem cell maintenance. Neuron, 114(2), 237.

Li H, et al. (2024) Bidirectional substrate shuttling between the 26S proteasome and the Cdc48 ATPase promotes protein degradation. Molecular cell.

Thornburg-Suresh EJC, et al. (2023) The Stathmin-2 membrane targeting domain is required for axon protection and regulated degradation by DLK signaling. The Journal of biological chemistry, 104861.

Radenkovic S, et al. (2023) Tracer metabolomics reveals the role of aldose reductase in glycosylation. Cell reports. Medicine, 4(6), 101056.

Rauckhorst AJ, et al. (2022) Mouse tissue harvest-induced hypoxia rapidly alters the in vivo metabolome, between-genotype metabolite level differences, and 13C-tracing enrichments. Molecular metabolism, 66, 101596.

Sharma A, et al. (2019) Impaired skeletal muscle mitochondrial pyruvate uptake rewires glucose metabolism to drive whole-body leanness. eLife, 8.