

Resource Summary Report

Generated by ASWG on Aug 10, 2026

Myeloid marker [BM2] antibody (ready-to-use)

RRID:AB_374280

Type: Antibody

Proper Citation

GeneTex Cat# GTX20908, RRID:AB_374280

Antibody Information

URL: http://antibodyregistry.org/AB_374280

Proper Citation: GeneTex Cat# GTX20908, RRID:AB_374280

Target Antigen: Myeloid marker [BM2] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.

Antibody Name: Myeloid marker [BM2] antibody (ready-to-use)

Description: This monoclonal targets Myeloid marker [BM2] antibody (ready-to-use)

Antibody ID: AB_374280

Vendor: GeneTex

Catalog Number: GTX20908

Record Creation Time: 20250819T232249+0000

Record Last Update: 20250820T051024+0000

Ratings and Alerts

No rating or validation information has been found for Myeloid marker [BM2] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.