

Resource Summary Report

Generated by [nidm-terms](#) on Aug 16, 2026

Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP

RRID:AB_325994

Type: Antibody

Proper Citation

Thermo Fisher Scientific Cat# SA1-200, RRID:AB_325994

Antibody Information

URL: http://antibodyregistry.org/AB_325994

Proper Citation: Thermo Fisher Scientific Cat# SA1-200, RRID:AB_325994

Target Antigen: Rabbit IgG (H+L)

Host Organism: donkey

Clonality: polyclonal secondary

Comments: Applications: WB (1:60-1:5,000)

Antibody Name: Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP

Description: This polyclonal secondary targets Rabbit IgG (H+L)

Target Organism: rabbit

Defining Citation: [PMID:24777581](#), [PMID:23794138](#), [PMID:14704233](#), [PMID:24470146](#)

Antibody ID: AB_325994

Vendor: Thermo Fisher Scientific

Catalog Number: SA1-200

Record Creation Time: 20250820T035017+0000

Record Last Update: 20250820T171939+0000

Ratings and Alerts

No rating or validation information has been found for Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP.

No alerts have been found for Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 10 mentions in open access literature.

Listed below are recent publications. The full list is available at [nidm-terms](#) .

Raina A, et al. (2025) Skin-derived α -synuclein strains from PD, DLB, and MSA induce distinct intracellular pathology and neurodegeneration. The Journal of biological chemistry, 302(1), 111005.

Krach F, et al. (2025) RNA splicing modulator for Huntington's disease treatment induces peripheral neuropathy. iScience, 28(5), 112380.

Carpentier M, et al. (2025) Seipin Governs caveolin-1 trafficking through modulating sphingolipid-glycerolipid balance. Cell reports, 44(10), 116320.

Li B, et al. (2024) Neural stem cell-derived exosomes promote mitochondrial biogenesis and restore abnormal protein distribution in a mouse model of Alzheimer's disease. Neural regeneration research, 19(7), 1593.

Koster KP, et al. (2024) Akap5 links synaptic dysfunction to neuroinflammatory signaling in a mouse model of infantile neuronal ceroid lipofuscinosis. Frontiers in synaptic neuroscience, 16, 1384625.

Benwell CJ, et al. (2024) A proteomics approach to isolating neuropilin-dependent α 5 integrin trafficking pathways: neuropilin 1 and 2 co-traffic α 5 integrin through endosomal p120RasGAP to promote polarised fibronectin fibrillogenesis in endothelial cells. Communications biology, 7(1), 629.

Koster KP, et al. (2023) Loss of Depalmitoylation Disrupts Homeostatic Plasticity of AMPARs in a Mouse Model of Infantile Neuronal Ceroid Lipofuscinosis. The Journal of neuroscience : the official journal of the Society for Neuroscience, 43(49), 8317.

Yap L, et al. (2019) In Vivo Generation of Post-infarct Human Cardiac Muscle by Laminin-Promoted Cardiovascular Progenitors. Cell reports, 26(12), 3231.

Hofmann F, et al. (2019) Cx43-Gap Junctions Accumulate at the Cytotoxic Immunological Synapse Enabling Cytotoxic T Lymphocyte Melanoma Cell Killing. International journal of molecular sciences, 20(18).

Sim J, et al. (2018) The Factor Inhibiting HIF Asparaginyl Hydroxylase Regulates Oxidative Metabolism and Accelerates Metabolic Adaptation to Hypoxia. *Cell metabolism*, 27(4), 898.