

# Resource Summary Report

Generated by [nidm-terms](#) on Aug 10, 2026

## Human Albumin antibody

RRID:AB\_374553

Type: Antibody

---

### Proper Citation

GeneTex Cat# GTX28940, RRID:AB\_374553

---

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_374553](http://antibodyregistry.org/AB_374553)

**Proper Citation:** GeneTex Cat# GTX28940, RRID:AB\_374553

**Target Antigen:** Human Albumin antibody

**Host Organism:** sheep

**Clonality:** polyclonal

**Comments:** Discontinued; manufacturer recommendations: IgG; IgG Dot Blot; Immunodiffusion; Other; Western Blot; ELISA; Immunohistochemistry; RID and Rocket IEP: 2µL antiserum/cm<sup>2</sup> gel vs 5µL 1/200 - 1/2000 dilution human serumDouble diffusion: 10µL antiserum vs 5µL serum (1/20 dilution)IEP: 100µL antiserum vs 1µL serumThe use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Suitability for use in nephelometry, immunohistochemical procedures, enzyme-linked immunosorbentassays and Western blot has not been assessed but use in such assays should not necessarily beexcluded. Double immunodiffusion, Immunoelectrophoresis, Radial immunodiffusion, Rocket immunoelectrophoresis.

**Antibody Name:** Human Albumin antibody

**Description:** This polyclonal targets Human Albumin antibody

**Target Organism:** feline, rat, porcine, canine, cow, goat, pig, horse, mouse, chickenbird, rabbit, cat, bovine, human, dog

**Antibody ID:** AB\_374553

**Vendor:** GeneTex

**Catalog Number:** GTX28940

**Record Creation Time:** 20250820T000804+0000

## Ratings and Alerts

No rating or validation information has been found for Human Albumin antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Dot Blot; Immunodiffusion; Other; Western Blot; ELISA; Immunohistochemistry; RID and Rocket IEP: 2µL antiserum/cm<sup>2</sup> gel vs 5µL 1/200 - 1/2000 dilution human serumDouble diffusion: 10µL antiserum vs 5µL serum (1/20 dilution)IEP: 100µL antiserum vs 1µL serumThe use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Suitability for use in nephelometry, immunohistochemical procedures, enzyme-linked immunosorbentassays and Western blot has not been assessed but use in such assays should not necessarily beexcluded. Double immunodiffusion, Immunoelectrophoresis, Radial immunodiffusion, Rocket immunoelectrophoresis.

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.