Resource Summary Report

Generated by PRECISE-TBI on Aug 24, 2026

Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 635

RRID:AB_2536186
Type: Antibody

Proper Citation

(Thermo Fisher Scientific Cat# A-31576, RRID:AB_2536186)

Antibody Information

URL: http://antibodyregistry.org/AB_2536186

Proper Citation: (Thermo Fisher Scientific Cat# A-31576, RRID:AB_2536186)

Target Antigen: Rabbit IgG (H+L)

Host Organism: goat

Clonality: polyclonal secondary

Comments: Applications: Flow, ICC/IF, WB

Antibody Name: Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 635

Description: This polyclonal secondary targets Rabbit IgG (H+L)

Target Organism: rabbit

Defining Citation: PMID:20674607, PMID:21503146, PMID:23161523, PMID:21861635, PMID:24192075, PMID:22811676, PMID:21428721, PMID:21441982

Antibody ID: AB_2536186

Vendor: Thermo Fisher Scientific

Catalog Number: A-31576

Record Creation Time: 20251213T073628+0000

Record Last Update: 20260225T230802+0000

Ratings and Alerts

No rating or validation information has been found for Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 635.

No alerts have been found for Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 635.

Data and Source Information

Source: Antibody Registry

Usage and Citation Metrics

We found 9 mentions in open access literature.

Listed below are recent publications. The full list is available at PRECISE-TBI .

Xu Y, et al. (2025) The cGAS-STING pathway activates transcription factor TFEB to stimulate lysosome biogenesis and pathogen clearance. Immunity, 58(2), 309.

Rajan G, et al. (2022) Evolutionary divergence of locomotion in two related vertebrate species. Cell reports, 38(13), 110585.

Chen N, et al. (2022) Local translation provides the asymmetric distribution of CaMKII required for associative memory formation. Current biology : CB, 32(12), 2730.

Reinhard N, et al. (2022) The lateral posterior clock neurons of Drosophila melanogaster express three neuropeptides and have multiple connections within the circadian clock network and beyond. The Journal of comparative neurology, 530(9), 1507.

Chaouch A, et al. (2021) Histone H3.3 K27M and K36M mutations de-repress transposable elements through perturbation of antagonistic chromatin marks. Molecular cell, 81(23), 4876.

Na HJ, et al. (2020) Nutrient-Driven O-GlcNAcylation Controls DNA Damage Repair Signaling and Stem/Progenitor Cell Homeostasis. Cell reports, 31(6), 107632.

Liu C, et al. (2019) A Serotonin-Modulated Circuit Controls Sleep Architecture to Regulate Cognitive Function Independent of Total Sleep in Drosophila. Current biology : CB, 29(21), 3635.

Schubert FK, et al. (2018) Neuroanatomical details of the lateral neurons of Drosophila melanogaster support their functional role in the circadian system. The Journal of comparative neurology, 526(7), 1209.

Nilssen ES, et al. (2018) Inhibitory Connectivity Dominates the Fan Cell Network in Layer II of Lateral Entorhinal Cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience, 38(45), 9712.