

Resource Summary Report

Generated by PRECISE-TBI on Aug 13, 2026

IL7 antibody

RRID:AB_371949

Type: Antibody

Proper Citation

GeneTex Cat# GTX10766, RRID:AB_371949

Antibody Information

URL: http://antibodyregistry.org/AB_371949

Proper Citation: GeneTex Cat# GTX10766, RRID:AB_371949

Target Antigen: IL7 antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG ELISA: Use at a concentration of 0.5 - 1.0 µg/ml. Neut: Use at an assay dependent dilution. WB: Use at a concentration of 0.1 - 0.2 µg/ml. Detects a band of approximately 15 kDa. The Neutralization Dose 50 (ND50) for anti-mouse IL7 is approximately 0.2-0.4µg/ml in the presence of 1.5 ng/ml of recombinant mouse IL7 using human peripheral blood mononuclear cells activated with PHA for 5 days. The exact concentration of antibody required to neutralize the recombinant mouse IL7 activity is dependent on the cytokine concentration, cell type, growth conditions, and the type of activity. The detection limit in immunoblotting for recombinant mouse IL7 is approximately 5.0 ng/lane under non-reducing and reducing conditions. The detection limit in ELISA for recombinant mouse IL7 is

Antibody Name: IL7 antibody

Description: This polyclonal targets IL7 antibody

Target Organism: mouse

Antibody ID: AB_371949

Vendor: GeneTex

Catalog Number: GTX10766

Record Creation Time: 20250820T020647+0000

Record Last Update: 20250820T124326+0000

Ratings and Alerts

No rating or validation information has been found for IL7 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ELISA: Use at a concentration of 0.5 - 1.0 µg/ml. Neut: Use at an assay dependent dilution. WB: Use at a concentration of 0.1 - 0.2 µg/ml. Detects a band of approximately 15 kDa. The Neutralization Dose 50 (ND50) for anti-mouse IL7 is approximately 0.2-0.4µg/ml in the presence of 1.5 ng/ml of recombinant mouse IL7 using human peripheral blood mononuclear cells activated with PHA for 5 days. The exact concentration of antibody required to neutralize the recombinant mouse IL7 activity is dependent on the cytokine concentration, cell type, growth conditions, and the type of activity. The detection limit in immunoblotting for recombinant mouse IL7 is approximately 5.0 ng/lane under non-reducing and reducing conditions. The detection limit in ELISA for recombinant mouse IL7 is

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.