

Resource Summary Report

Generated by PRECISE-TBI on Aug 31, 2026

C1r antibody

RRID:AB_374514

Type: Antibody

Proper Citation

GeneTex Cat# GTX28781, RRID:AB_374514

Antibody Information

URL: http://antibodyregistry.org/AB_374514

Proper Citation: GeneTex Cat# GTX28781, RRID:AB_374514

Target Antigen: C1r antibody

Host Organism: sheep

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG ELISA; Other; Western Blot; Immunodiffusion; Immunohistochemistry; RID and Rocket IEP: 2µL antiserum/cm² in gel vs 20µL fresh plasma, NT - 1/2Double Diffusion: 10µL antiserum vs 20µL fresh plasmaSecond Antibody: 100µL antiserum vs 5µL fresh plasma The use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Results can be enhanced by washing completed agarose gels in saline, drying and finally stainingwith a protein stain such as coomassie blue. Suitability for use in immunohistochemical procedures, nephelometry, enzyme-linked immunosorbent assays and Western blot has not been assessed butuse in such assays should not necessarily be excluded. Double immunodiffusion, Immunoelectrophoresis, Radial imm

Antibody Name: C1r antibody

Description: This polyclonal targets C1r antibody

Target Organism: human

Antibody ID: AB_374514

Vendor: GeneTex

Catalog Number: GTX28781

Record Creation Time: 20250819T233123+0000

Ratings and Alerts

No rating or validation information has been found for C1r antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ELISA; Other; Western Blot; Immunodiffusion; Immunohistochemistry; RID and Rocket IEP: 2µL antiserum/cm² in gel vs 20µL fresh plasma, NT - 1/2Double Diffusion: 10µL antiserum vs 20µL fresh plasmaSecond Antibody: 100µL antiserum vs 5µL fresh plasma The use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Results can be enhanced by washing completed agarose gels in saline, drying and finally staining with a protein stain such as coomassie blue. Suitability for use in immunohistochemical procedures, nephelometry, enzyme-linked immunosorbent assays and Western blot has not been assessed but use in such assays should not necessarily be excluded. Double immunodiffusion, Immunoelectrophoresis, Radial imm

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.