

Resource Summary Report

Generated by [PRECISE-TBI](#) on Aug 9, 2026

c-Myc [CT14.G4] antibody

RRID:AB_377285

Type: Antibody

Proper Citation

GeneTex Cat# GTX76330, RRID:AB_377285

Antibody Information

URL: http://antibodyregistry.org/AB_377285

Proper Citation: GeneTex Cat# GTX76330, RRID:AB_377285

Target Antigen: c-Myc [CT14.G4] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow Cytometry; Immunofluorescence; Immunohistochemistry - fixed; Western Blot; Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - frozen; For FACS, IHC-Fr, IHC-P: Use at a dilution of 1:50 - 1:200. For IP and WB: Use at an assay dependent dilution. Not tested in other applications. Optimal concentrations/dilutions should be determined by the researchers. Immunohistology: ** This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH6.0 is recommended for this purpose., Flow cytometry/FACS, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-Fr, IHC-P, IP, WB

Antibody Name: c-Myc [CT14.G4] antibody

Description: This monoclonal targets c-Myc [CT14.G4] antibody

Target Organism: human

Antibody ID: AB_377285

Vendor: GeneTex

Catalog Number: GTX76330

Record Creation Time: 20250820T041444+0000

Record Last Update: 20250820T182640+0000

Ratings and Alerts

No rating or validation information has been found for c-Myc [CT14.G4] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow Cytometry; Immunofluorescence; Immunohistochemistry - fixed; Western Blot; Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - frozen; For FACS, IHC-Fr, IHC-P: Use at a dilution of 1:50 - 1:200. For IP and WB: Use at an assay dependent dilution. Not tested in other applications. Optimal concentrations/dilutions should be determined by the researchers. Immunohistology: ** This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH6.0 is recommended for this purpose., Flow cytometry/FACS, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-Fr, IHC-P, IP, WB

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.