

Resource Summary Report

Generated by PRECISE-TBI on Sep 3, 2026

Tubulin antibody [B3]

RRID:AB_381313

Type: Antibody

Proper Citation

GeneTex Cat# GTX11324, RRID:AB_381313

Antibody Information

URL: http://antibodyregistry.org/AB_381313

Proper Citation: GeneTex Cat# GTX11324, RRID:AB_381313

Target Antigen: Tubulin antibody [B3]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgM; IgM Immunocytochemistry; ELISA; Western Blot; ELISA, Immunocytochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, ICC, WB, ELISA: Use at an assay dependent dilution. ICC: Use at an assay dependent dilution. Can be used with 4 paraformaldehyde-1 Triton X-100 and methanol-acetone. WB: Use at a concentration of 0.5 - 1 µg/ml. This concentration is recommended using a cytosolic fraction of rat brain extract. Detects a band of approximately 50 kDa, possibly a doublet of alpha and beta tubulins. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: Tubulin antibody [B3]

Description: This monoclonal targets Tubulin antibody [B3]

Target Organism: chicken, rat, cow, mouse, chickenbird, rabbit, bovine, other invertebrate, human

Antibody ID: AB_381313

Vendor: GeneTex

Catalog Number: GTX11324

Record Creation Time: 20231110T080840+0000

Ratings and Alerts

No rating or validation information has been found for Tubulin antibody [B3].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgM; IgM Immunocytochemistry; ELISA; Western Blot; ELISA, Immunocytochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, ICC, WB, ELISA: Use at an assay dependent dilution. ICC: Use at an assay dependent dilution. Can be used with 4 paraformaldehyde-1 Triton X-100 and methanol-acetone. WB: Use at a concentration of 0.5 - 1 µg/ml. This concentration is recommended using a cytosolic fraction of rat brain extract. Detects a band of approximately 50 kDa, possibly a doublet of alpha and beta tubulins. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 2 mentions in open access literature.

Listed below are recent publications. The full list is available at [PRECISE-TBI](#) .

Aluru K, et al. (2025) Protocol for purifying primitive neural progenitors by fractional passaging to generate astroglia-enriched cerebral organoid. STAR protocols, 6(4), 104142.

Kim H, et al. (2021) Generation of human pluripotent stem cell-derived fused organoids with oligodendroglia and myelin. STAR protocols, 2(2), 100443.