

Resource Summary Report

Generated by SPARC SAWG on Sep 3, 2026

Myelin basic protein [B103.1] antibody

RRID:AB_1080897

Type: Antibody

Proper Citation

GeneTex Cat# GTX72695, RRID:AB_1080897

Antibody Information

URL: http://antibodyregistry.org/AB_1080897

Proper Citation: GeneTex Cat# GTX72695, RRID:AB_1080897

Target Antigen: Myelin basic protein [B103.1] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody can be used on formalin-fixed, paraffin-embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of 1:25 to 1:50 with AutoProbe III (GTX28620). When using paraffin-embedded and formalin-fixed tissues, high temperature antigenic unmasking technique (see instructions below) is strongly recommended for consistent and reproducible results. This product can also be used on frozen tissue sections. The optimal conditions should be determined by the individual laborat

Antibody Name: Myelin basic protein [B103.1] antibody

Description: This monoclonal targets Myelin basic protein [B103.1] antibody

Target Organism: monkey, porcine, chickenbird, rabbit, bovine, human

Antibody ID: AB_1080897

Vendor: GeneTex

Catalog Number: GTX72695

Record Creation Time: 20231110T060333+0000

Record Last Update: 20260829T141443+0000

Ratings and Alerts

No rating or validation information has been found for Myelin basic protein [B103.1] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody can be used on formalin-fixed, paraffin-embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of 1:25 to 1:50 with AutoProbe III (GTX28620). When using paraffin-embedded and formalin-fixed tissues, high temperature antigenic unmasking technique (see instructions below) is strongly recommended for consistent and reproducible results. This product can also be used on frozen tissue sections. The optimal conditions should be determined by the individual laborat

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.