

Resource Summary Report

Generated by SPARC SAWG on Aug 11, 2026

TNF alpha antibody

RRID:AB_367845

Type: Antibody

Proper Citation

GeneTex Cat# GTX10949, RRID:AB_367845

Antibody Information

URL: http://antibodyregistry.org/AB_367845

Proper Citation: GeneTex Cat# GTX10949, RRID:AB_367845

Target Antigen: TNF alpha antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; ELISA, IHC (Formalin-fixed paraffin-embedded sections), Neutralizing, Western blot. For ELISA: Use at a concentration of 0.5 - 1 µg/ml to detect 0.06 ng/well of rhTNF alpha. For IHC-P: Use at a concentration of 0.5 - 5 µg/ml. For Neut: Use at a concentration of 0.003 - 0.01 µg/ml. For WB: Use at a concentration of 0.1 - 0.2 µg/ml to detect rhTNF alpha at 0.2 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 17 kDa. Anti-Human TNF alpha is tested for its ability to neutralize the bioactivity of rhTNF-alpha in a cytotoxicity assay using murine L-929. The ND50 of the antibody is defined as the concentration of antibody resulting in a one-h

Antibody Name: TNF alpha antibody

Description: This polyclonal targets TNF alpha antibody

Target Organism: rat, mouse, human

Antibody ID: AB_367845

Vendor: GeneTex

Catalog Number: GTX10949

Record Creation Time: 20250819T233248+0000

Ratings and Alerts

No rating or validation information has been found for TNF alpha antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; ELISA, IHC (Formalin-fixed paraffin-embedded sections), Neutralizing, Western blot. For ELISA: Use at a concentration of 0.5 - 1 µg/ml to detect 0.06 ng/well of rhTNF alpha. For IHC-P: Use at a concentration of 0.5 - 5 µg/ml. For Neut: Use at a concentration of 0.003 - 0.01 µg/ml. For WB: Use at a concentration of 0.1 - 0.2 µg/ml to detect rhTNF alpha at 0.2 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 17 kDa. Anti-Human TNF alpha is tested for its ability to neutralize the bioactivity of rhTNF-alpha in a cytotoxicity assay using murine L-929. The ND50 of the antibody is defined as the concentration of antibody resulting in a one-h

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.