

Resource Summary Report

Generated by SPARC SAWG on Aug 18, 2026

Clathrin Light chain [3F133] antibody

RRID:AB_369624

Type: Antibody

Proper Citation

GeneTex Cat# GTX14408, RRID:AB_369624

Antibody Information

URL: http://antibodyregistry.org/AB_369624

Proper Citation: GeneTex Cat# GTX14408, RRID:AB_369624

Target Antigen: Clathrin Light chain [3F133] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunofluorescence; Western Blot; Immunohistochemistry - fixed; Immunohistochemistry; Immunoprecipitation; IF, IHC-P, IP, WB, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IF: Use at an assay dependent dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml for 30 min at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution with protein A. WB: Use at a concentration of 1 - 2 µg/ml for 2hrs at RT. Predicted molecular weight: 28 kDa. Not tested in other applications. Optimal dilutions/concentrations should b

Antibody Name: Clathrin Light chain [3F133] antibody

Description: This monoclonal targets Clathrin Light chain [3F133] antibody

Target Organism: rat, xenopusamphibian, mouse, zebrafishfish, bovine, human

Antibody ID: AB_369624

Vendor: GeneTex

Catalog Number: GTX14408

Record Creation Time: 20250819T211746+0000

Ratings and Alerts

No rating or validation information has been found for Clathrin Light chain [3F133] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunofluorescence; Western Blot; Immunohistochemistry - fixed; Immunohistochemistry; Immunoprecipitation; IF, IHC-P, IP, WB, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IF: Use at an assay dependent dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml for 30 min at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution with protein A. WB: Use at a concentration of 1 - 2 µg/ml for 2hrs at RT. Predicted molecular weight: 28 kDa. Not tested in other applications. Optimal dilutions/concentrations should b

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.