

Resource Summary Report

Generated by SPARC SAWG on Aug 10, 2026

Eotaxin 2 antibody [61016.11]

RRID:AB_373980

Type: Antibody

Proper Citation

GeneTex Cat# GTX10370, RRID:AB_373980

Antibody Information

URL: http://antibodyregistry.org/AB_373980

Proper Citation: GeneTex Cat# GTX10370, RRID:AB_373980

Target Antigen: Eotaxin 2 antibody [61016.11]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 ELISA, Neutralizing, Western blot. The usefulness of this product in other applications has not been determined., ELISA, Neut, WB, ELISA: Use at a concentration of 2 µg/ml. Neut: Use at a concentration of 1 - 5 µg/ml. WB: Use at a concentration of 1 - 2 µg/ml. Predicted molecular weight: 10.6 kDa. The detection limit for human eotaxin 2 is approximately 1 ng/lane under non-reducing and reducing conditions. Eotaxin 2 has the ability to neutralize the biological activity of recombinant human eotaxin 2. To measure this activity, recombinant human eotaxin 2 is incubated with various concentrations of the antibody for 30 minutes at room temperature in a 96 well microplate. Following this preincubation period, 35 µl of the cytokine-antibody solution (containing recombinant

Antibody Name: Eotaxin 2 antibody [61016.11]

Description: This monoclonal targets Eotaxin 2 antibody [61016.11]

Target Organism: human

Antibody ID: AB_373980

Vendor: GeneTex

Catalog Number: GTX10370

Record Creation Time: 20250820T001218+0000

Ratings and Alerts

No rating or validation information has been found for Eotaxin 2 antibody [61016.11].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 ELISA, Neutralizing, Western blot. The usefulness of this product in other applications has not been determined., ELISA, Neut, WB, ELISA: Use at a concentration of 2 µg/ml. Neut: Use at a concentration of 1 - 5 µg/ml. WB: Use at a concentration of 1 - 2 µg/ml. Predicted molecular weight: 10.6 kDa. The detection limit for human eotaxin 2 is approximately 1 ng/lane under non-reducing and reducing conditions. Eotaxin 2 has the ability to neutralizethe biological activity of recombinant human eotaxin 2. To measure this activity, recombinant human eotaxin 2 is incubated with various concentrations of the antibody for 30 minutes at room temperature in a 96 well microplate. Following this preincubation period, 35 µl of the cytokine-antibody solution (containing recombinant

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.