

Resource Summary Report

Generated by SPARC SAWG on Sep 5, 2026

Retinoid X Receptor beta antibody [MOK13-17]

RRID:AB_384870

Type: Antibody

Proper Citation

GeneTex Cat# GTX22815, RRID:AB_384870

Antibody Information

URL: http://antibodyregistry.org/AB_384870

Proper Citation: GeneTex Cat# GTX22815, RRID:AB_384870

Target Antigen: Retinoid X Receptor beta antibody [MOK13-17]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Immunofluorescence; Immunocytochemistry; Immunoprecipitation; Western Blot; Super Shift Assay; Gel supershift assays, Immunocytochemistry, Immunofluorescence, Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., GSA, ICC, IF, IP, WB, For WB: use at a concentration of 5 µg/mlGel Shift: Assay dependent. For ICC/IF: Assay dependent. For IP: Assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects an ~50 kDa protein representing RXR beta from transfected Sf9 cells. For ICC/IF staining of RXR beta in MCF-7 cells overexpressing RXR beta results in strong nuclear staining. Gel super shift experiments show disruption of vitamin

Antibody Name: Retinoid X Receptor beta antibody [MOK13-17]

Description: This monoclonal targets Retinoid X Receptor beta antibody [MOK13-17]

Target Organism: mouse, human

Antibody ID: AB_384870

Vendor: GeneTex

Catalog Number: GTX22815

Record Creation Time: 20231110T060136+0000

Record Last Update: 20260829T055122+0000

Ratings and Alerts

No rating or validation information has been found for Retinoid X Receptor beta antibody [MOK13-17].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Immunofluorescence; Immunocytochemistry; Immunoprecipitation; Western Blot; Super Shift Assay; Gel supershift assays, Immunocytochemistry, Immunofluorescence, Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., GSA, ICC, IF, IP, WB, For WB: use at a concentration of 5 µg/mlGel Shift: Assay dependent. For ICC/IF: Assay dependent. For IP: Assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects an ~50 kDa protein representing RXR beta from transfected Sf9 cells. For ICC/IF staining of RXR beta in MCF-7 cells overexpressing RXR beta results in strong nuclear staining. Gel super shift experiments show disruption of vitamin

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.