

Resource Summary Report

Generated by SPARC SAWG on Sep 16, 2026

DMLE

RRID:SCR_013454

Type: Tool

Proper Citation

DMLE (RRID:SCR_013454)

Resource Information

URL: <http://www.dmle.org/>

Proper Citation: DMLE (RRID:SCR_013454)

Description: Software application for high-resolution mapping of the position of a disease mutation relative to a set of genetic markers using population linkage disequilibrium (LD). (entry from Genetic Analysis Software)

Abbreviations: DMLE

Synonyms: Disease Mapping using Linkage disEquilibrium

Resource Type: software application, software resource

Keywords: gene, genetic, genomic, linux, ms-windows

Resource Name: DMLE

Resource ID: SCR_013454

Alternate IDs: nlx_154218

Record Creation Time: 20220129T080316+0000

Record Last Update: 20260912T200249+0000

Ratings and Alerts

No rating or validation information has been found for DMLE.

No alerts have been found for DMLE.

Data and Source Information

Usage and Citation Metrics

We found 22 mentions in open access literature.

Listed below are recent publications. The full list is available at [SPARC SAWG](#) .

Danilchenko VY, et al. (2025) Comparative analysis of haplotypes carrying pathogenic variants c.1545T>G, c.2027T>A and c.919-2A>G of the SLC26A4 gene in patients with hearing loss from the Tyva Republic (Southern Siberia). Vavilovskii zhurnal genetiki i selektsii, 29(1), 144.

Danilchenko VY, et al. (2023) Insight into the Natural History of Pathogenic Variant c.919-2A>G in the SLC26A4 Gene Involved in Hearing Loss: The Evidence for Its Common Origin in Southern Siberia (Russia). Genes, 14(4).

Islam S, et al. (2021) Founder mutation in the PMM2 promotor causes hyperinsulinemic hypoglycaemia/polycystic kidney disease (HIPKD). Molecular genetics & genomic medicine, 9(12), e1674.

Luo X, et al. (2021) Characterization of a Cohort of Patients With LIG4 Deficiency Reveals the Founder Effect of p.R278L, Unique to the Chinese Population. Frontiers in immunology, 12, 695993.

Zytsar MV, et al. (2020) High Rates of Three Common GJB2 Mutations c.516G>C, c.-23+1G>A, c.235delC in Deaf Patients from Southern Siberia Are Due to the Founder Effect. Genes, 11(7).

Kim YM, et al. (2020) The GBA p.G85E mutation in Korean patients with non-neuronopathic Gaucher disease: founder and neuroprotective effects. Orphanet journal of rare diseases, 15(1), 318.

Esperón-Moldes U, et al. (2020) Novel CYP4F22 mutations associated with autosomal recessive congenital ichthyosis (ARCI). Study of the CYP4F22 c.1303C>T founder mutation. PloS one, 15(2), e0229025.

Tatour Y, et al. (2019) A novel intronic mutation of PDE6B is a major cause of autosomal recessive retinitis pigmentosa among Caucasus Jews. Molecular vision, 25, 155.

Garagiola I, et al. (2016) A recurrent F8 mutation (c.6046C>T) causing hemophilia A in 8% of northern Italian patients: evidence for a founder effect. Molecular genetics & genomic medicine, 4(2), 152.

Caleca L, et al. (2014) Characterization of an Italian founder mutation in the RING-finger domain of BRCA1. PloS one, 9(2), e86924.

de Alencar DO, et al. (2014) Fabry disease: Evidence for a regional founder effect of the GLA gene mutation 30delG in Brazilian patients. Molecular genetics and metabolism reports, 1, 414.

Winbo A, et al. (2014) Phenotype, origin and estimated prevalence of a common long QT syndrome mutation: a clinical, genealogical and molecular genetics study including Swedish R518X/KCNQ1 families. BMC cardiovascular disorders, 14, 22.

García-Murias M, et al. (2012) 'Costa da Morte' ataxia is spinocerebellar ataxia 36: clinical and genetic characterization. Brain : a journal of neurology, 135(Pt 5), 1423.

Liskova P, et al. (2012) High prevalence of posterior polymorphous corneal dystrophy in the Czech Republic; linkage disequilibrium mapping and dating an ancestral mutation. PloS one, 7(9), e45495.

Hünemeier T, et al. (2012) Evolutionary responses to a constructed niche: ancient Mesoamericans as a model of gene-culture coevolution. PloS one, 7(6), e38862.

Fachal L, et al. (2012) Multiple local and recent founder effects of TGM1 in Spanish families. PloS one, 7(4), e33580.

Borroni B, et al. (2011) Founder effect and estimation of the age of the Progranulin Thr272fs mutation in 14 Italian pedigrees with frontotemporal lobar degeneration. Neurobiology of aging, 32(3), 555.e1.

Semmler A, et al. (2011) Haplotype analysis of the 5,10-methylenetetrahydrofolate reductase (MTHFR) c.1298A>C (E429A) polymorphism. BMC research notes, 4, 439.

Cornes BK, et al. (2010) Haplotype analysis reveals a possible founder effect of RET mutation R114H for Hirschsprung's disease in the Chinese population. PloS one, 5(6), e10918.

Cichon S, et al. (2006) Increased activity of coagulation factor XII (Hageman factor) causes hereditary angioedema type III. American journal of human genetics, 79(6), 1098.