

Resource Summary Report

Generated by [FDI Lab - SciCrunch.org](#) on May 22, 2025

Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Cyanine3

RRID:AB_10563288

Type: Antibody

Proper Citation

(Thermo Fisher Scientific Cat# A10520, RRID:AB_10563288)

Antibody Information

URL: http://antibodyregistry.org/AB_10563288

Proper Citation: (Thermo Fisher Scientific Cat# A10520, RRID:AB_10563288)

Target Antigen: Rabbit IgG (H+L)

Host Organism: goat

Clonality: polyclonal secondary

Comments: Applications: ICC/IF (1-10 µg/mL), WB (1:10,000)

Antibody Name: Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Cyanine3

Description: This polyclonal secondary targets Rabbit IgG (H+L)

Target Organism: rabbit

Defining Citation: [PMID:27821182](#), [PMID:27068301](#), [PMID:27698380](#), [PMID:23603844](#)

Antibody ID: AB_10563288

Vendor: Thermo Fisher Scientific

Catalog Number: A10520

Record Creation Time: 20241130T060402+0000

Ratings and Alerts

No rating or validation information has been found for Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Cyanine3.

No alerts have been found for Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Cyanine3.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 12 mentions in open access literature.

Listed below are recent publications. The full list is available at [FDI Lab - SciCrunch.org](#).

Zhang Y, et al. (2025) Gamma-glutamyl transferase 5 overexpression in cerebrovascular endothelial cells improves brain pathology, cognition, and behavior in APP/PS1 mice. *Neural regeneration research*, 20(2), 533.

Concina G, et al. (2024) Hippocampus-to-amygdala pathway drives the separation of remote memories of related events. *Cell reports*, 43(5), 114151.

De Gasperi R, et al. (2024) Septin 7 interacts with Numb to preserve sarcomere structural organization and muscle contractile function. *eLife*, 12.

Carabaña C, et al. (2024) Spatially distinct epithelial and mesenchymal cell subsets along progressive lineage restriction in the branching embryonic mammary gland. *The EMBO journal*, 43(12), 2308.

Onji H, et al. (2024) Schlafen 11 further sensitizes BRCA-deficient cells to PARP inhibitors through single-strand DNA gap accumulation behind replication forks. *Oncogene*, 43(32), 2475.

Vaughen JP, et al. (2022) Glial control of sphingolipid levels sculpts diurnal remodeling in a circadian circuit. *Neuron*, 110(19), 3186.

Foray V, et al. (2018) Wolbachia Control Stem Cell Behavior and Stimulate Germline Proliferation in Filarial Nematodes. *Developmental cell*, 45(2), 198.

Frey E, et al. (2018) TRPV1 Agonist, Capsaicin, Induces Axon Outgrowth after Injury via Ca²⁺/PKA Signaling. *eNeuro*, 5(3).

Kaptan E, et al. (2017) Runt-Related Transcription Factor 2 (Runx2) Is Responsible for Galectin-3 Overexpression in Human Thyroid Carcinoma. *Journal of cellular biochemistry*, 118(11), 3911.

Takagi S, et al. (2017) Divergent Connectivity of Homologous Command-like Neurons Mediates Segment-Specific Touch Responses in *Drosophila*. *Neuron*, 96(6), 1373.

Schmöle AC, et al. (2015) Expression Analysis of CB2-GFP BAC Transgenic Mice. *PloS one*, 10(9), e0138986.

Schmöle AC, et al. (2015) Cannabinoid receptor 2 deficiency results in reduced neuroinflammation in an Alzheimer's disease mouse model. *Neurobiology of aging*, 36(2), 710.