Resource Summary Report

Generated by FDI Lab - SciCrunch.org on May 7, 2025

EGF antibody [EGF-10]

RRID:AB_381029 Type: Antibody

Proper Citation

(GeneTex Cat# GTX10409, RRID:AB_381029)

Antibody Information

URL: http://antibodyregistry.org/AB_381029

Proper Citation: (GeneTex Cat# GTX10409, RRID:AB_381029)

Target Antigen: EGF antibody [EGF-10]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Immunohistochemistry - fixed; Radioimmunoassay; Western Blot; Dot Blot; Immunohistochemistry; ELISA; Dot blot, ELISA, IHC (Formalin-fixed paraffin-embedded sections), Radioimmunoassay, Western blot. The usefulness of this product in other applications has not been determined., Dot, ELISA, IHC-P, RIA, WB, Dot: Use at a concentration of 1 μg/ml. ELISA: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. RIA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 144 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: EGF antibody [EGF-10]

Description: This monoclonal targets EGF antibody [EGF-10]

Target Organism: human

Antibody ID: AB 381029

Vendor: GeneTex

Catalog Number: GTX10409

Record Creation Time: 20241016T232854+0000

Record Last Update: 20241017T004532+0000

Ratings and Alerts

No rating or validation information has been found for EGF antibody [EGF-10].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Immunohistochemistry - fixed; Radioimmunoassay; Western Blot; Dot Blot; Immunohistochemistry; ELISA; Dot blot, ELISA, IHC (Formalin-fixed paraffin-embedded sections), Radioimmunoassay, Western blot. The usefulness of this product in other applications has not been determined., Dot, ELISA, IHC-P, RIA, WB, Dot: Use at a concentration of 1 µg/ml. ELISA: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. RIA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 144 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: Antibody Registry

Usage and Citation Metrics

We have not found any literature mentions for this resource.